



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

change the coefficient of  $a^2y^2$  from 96 to 100 the equation represents the straight lines  $y = \pm x$  and the circle  $x^2 + y^2 = 100$ , to which system of lines the curve is therefore closely asymptotic. I conjecture that the curve was originated in this way and has not heretofore been known as a geometrical locus. Can any of your readers throw any light on the history of this curious curve and its startling title?

---

ANSWER TO PROF. HALL'S QUERY, BY PROF. W. W. JOHNSON.

Prof. Cayley proposed the question "Find the number of regions into which infinite space is divided by  $n$  planes" in the Smith Prize Examination Feb. 3rd, 1874, and published in the Mathematical Messenger for march 1874, "Solutions and remarks" on the paper of that day. He says he intended the question for a problem, as the result, though a known, is not a generally known one. His solution is substantially as follows: Consider the analogous problem for lines in a plane. An additional line adds to the number of regions one for every part into which it is itself divided by the other lines. Hence, 1, 2, 3, 4 &c. lines divide a plane into 2,  $2+2(=4)$ ,  $4+3(=7)$ ,  $7+4(=11)$  &c. regions; the general term being  $\frac{1}{2}(n^2+n+2)$ . In like manner an additional plane adds to the number of regions in space one for every region in to which it is itself divided by the other planes. Hence 1, 2, 3, 4, &c. planes divide space into 2,  $2+2(=4)$ ,  $4+4(=8)$ ,  $8+7(=15)$ ,  $15+11(=26)$  &c. regions; the general term being  $\frac{1}{6}(n^3+5n+6)$ .

[Mr. G. W. Hill obtains the same result as answer to Prof. Hall's Query and by analogous reasoning, employing however in his investigation the Calculus of Finite Differences.

It will be observed that the question as proposed by Prof. Cayley is not identical with that proposed by Prof. Hall; as Prof. Cayley requests the number of regions into which infinite space is divided by  $n$  planes, whereas Prof. Hall asks, "Into how many parts *can*  $n$  planes divide space."

That the answer given is not *necessarily* the answer to Prof. Cayley's question follows from the fact that nothing in Prof. Cayley's announcement of the question precludes the possibility (theoretically at least) of some or all of the planes being parallel, in which case the answer would obviously not be correct: If drawn at random, however, the probability of such a contingency is infinitely small.—Ed.]

---

ANSWER TO PRESIDENT TAPPAN'S QUERY, BY PROF. J. SCHEFFER.

It is. The French mathematician *Fermat* who published quite a number of theorems in regard to prime numbers, erroneously asserted that all the